

**P4P Quality Measures
2007- 2008**

Speciality	Measurement	Description	Data Source
Orthopedic Surgery			
	Timing of Prophylactic Antibiotics Ordering Physician	Percentage of surgical patients aged 18 years and older undergoing procedures with the indications for prophylactic parenteral antibiotics who have an order for an antibiotic to be given within one hour (if vancomycin, two hours) prior to the surgical incision (or start of procedure when no incision is required)	MSS
	Timing of Prophylactic Antibiotics Administering Physician	Percentage of surgical patients aged 18 years and older who have an order for a parenteral antibiotic to be given within one hour (if vancomycin, two hours) prior to the surgical incision (or start of procedure when no incision is required) for whom administration of a prophylactic antibiotic has been initiated within one hour (if vancomycin, two hours) prior to the surgical incision (or start of procedure when no incision is required)	MSS
	Discontinuation of Prophylactic Antibiotics	Percentage of surgical patients aged 18 years and older undergoing procedures with the indications for prophylactic antibiotics AND who received a prophylactic antibiotic, who have an order for discontinuation of prophylactic antibiotics within 24 hours of surgical end time	MSS
	Deep Venous Thromboembolism (DVTE) Prophylaxis	Percentage of patients undergoing procedures for which DVTE prophylaxis is indicated in all patients who had an order for LMWH, LDUH, adjusted-dose warfarin, fondaparinux or mechanical prophylaxis to be given within 24 hours prior to incision time or 24 hours after surgery end time	MSS

**P4P Quality Measures
2007-2008**

Specialty	Measurement	Description	Source
Orthopedic Surgery	Complication Rate		MSS
	Re-admission Rate		MSS
	Mortality rate		MSS
	ALOS		
	Generic Drug Utilization	Percentage of generic drugs utilized	C
			MSS